

Study of a self-adjoint operator indicating the direction of time within standard quantum mechanics

Y. Strauss¹, J. Silman^{2*}, S. Machnes^{2†}, L.P. Horwitz^{3, 4}

¹*Einstein Institute of Mathematics, Edmond J. Safra campus,*

The Hebrew University of Jerusalem, Jerusalem 91904, Israel

²*School of Physics and Astronomy, Raymond and Beverly Sackler*

Faculty of Exact Sciences, Tel-Aviv University, Tel-Aviv 69978, Israel

³*Physics Department of Physics, Bar-Ilan University, Ramat-Gan 52900, Israel*

⁴*Department of Physics, The Ariel University Center of Samaria, Ariel 40700, Israel*

In [J. Math. Phys. 51 (2010) 022104] a self-adjoint operator was introduced that has the property that it indicates the direction of time within the framework of standard quantum mechanics, in the sense that as a function of time its expectation value decreases monotonically for any initial state. In this paper we study some of this operator's properties. In particular, we derive its spectrum and generalized eigenstates, and treat the example of the free particle.

I. INTRODUCTION

It is a fundamental question in standard quantum mechanics (SQM) of what type of restrictions the Schrödinger evolution imposes on the behavior in time of basic objects. In particular, it is of interest to ask if SQM allows for self-adjoint operators having the so-called Lyapunov property, that is, monotonicity of the expectation value irrespective of the initial state of the system. Clearly, such an operator would indicate the direction of time.

A natural candidate for a self-adjoint Lyapunov operator is a (self-adjoint) time operator T canonically conjugate to the Hamiltonian H , such that T and H form an imprimitivity system [1] (implying that each generates a translation on the spectrum of the other). However, a well known theorem of Pauli tells us that this is impossible [2]. Recently, Galapon attempted to bypass Pauli's arguments and found pairs of T and H satisfying the canonical commutation relations, but do not constitute an imprimitivity system [3]. It can be shown that the T operator obtained in this way does not have the Lyapunov property. Other authors do not insist on the conjugacy of T and H . In this context, Unruh and Wald's proof that a 'monotonically perfect clock' does not exist [4] should be noted, as well as Misra, Prigogine, and Courbage's no-go theorem [5]. Still another solution, advocated by some authors, is to do away with the requirement of self-adjointness [6].

A Lyapunov self-adjoint operator acting within the framework of standard quantum mechanics was recently introduced in [7]. In this paper we study some of its properties. In particular, we derive its spectrum and generalized eigenstates, and treat the example of the free particle.

II. MAIN RESULTS

Let $\mathcal{H}$ be a separable Hilbert space and let H be a self-adjoint operator generating a unitary evolution group $\{U(t)\}_{t \in \mathbb{R}}$, with $U(t) = \exp(-iHt)$, on $\mathcal{H}$. We take $\mathcal{H}$ to represent the Hilbert space corresponding to some given quantum system and H its Hamiltonian. For an initial state $\psi(0) = \psi \in \mathcal{H}$ $\psi(t) = U(t)\psi$ denotes the state of the system at time t and $\Psi_\psi := \{\psi(t)\}_{t \in \mathbb{R}^+}$ its trajectory. Let $\mathcal{B}(\mathcal{H})$ be the space of bounded linear operators on $\mathcal{H}$. A (forward) Lyapunov operator is defined as follows [7]:

Definition II.1 *Let $M \in \mathcal{B}(\mathcal{H})$ be a self-adjoint operator on $\mathcal{H}$. Let Ψ_ψ be the trajectory corresponding to an initial state ψ . Let $M(\Psi_\psi) = \{\|\varphi\|^{-2}(\varphi, M\varphi) \mid \varphi \in \Psi_\psi\}$ be the set of expectation values of M in states in Ψ_ψ . Then M is a forward Lyapunov operator if the mapping $\tau_{M, \psi} : \mathbb{R}^+ \mapsto M(\Psi_\psi)$ defined by*

$$\tau_{M, \psi}(t) = \|\psi(t)\|^{-2}(\psi(t), M\psi(t)) = \|\psi\|^{-2}(\psi(t), M\psi(t)) \quad (1)$$

is one to one and monotonically decreasing for all non-recurring trajectories. □

* Present address: Laboratoire d'Information Quantique, Université Libre de Bruxelles, 1050 Bruxelles, Belgium

† Present address: Institut für Theoretische Physik, Universität Ulm, 89069 Ulm, Germany

Let $\mathcal{K}$ be a separable Hilbert space and $L^2(\mathbb{R}^+; \mathcal{K})$ the Hilbert space of Lebesgue square-integrable $\mathcal{K}$ valued functions defined on $\mathbb{R}^+$, let H be the operator of multiplication by the independent variable on $L^2(\mathbb{R}^+; \mathcal{K})$, and let $\{U(t)\}_{t \in \mathbb{R}}$ be the continuous one-parameter unitary evolution group generated by H , i.e.,

$$[U(t)f](E) = [e^{-iHt}f](E) = e^{-iEt}f(E), \quad f \in L^2(\mathbb{R}^+; \mathcal{K}), \quad E \in \mathbb{R}^+. \quad (2)$$

Denote by $\mathcal{H}^2(\mathbb{C}^\pm; \mathcal{K})$ the Hardy spaces of $\mathcal{K}$ valued functions analytic in $\mathbb{C}^\pm$. The Hilbert spaces $\mathcal{H}_\pm^2(\mathbb{R}; \mathcal{K})$ consisting of non-tangential boundary values on $\mathbb{R}$ in $\mathcal{H}^2(\mathbb{C}^\pm; \mathcal{K})$ are, respectively, isomorphic to $\mathcal{H}^2(\mathbb{C}^\pm; \mathcal{K})$. The spaces $\mathcal{H}_\pm^2(\mathbb{R}; \mathcal{K})$ are orthogonal subspaces of $L^2(\mathbb{R}; \mathcal{K})$ with $L^2(\mathbb{R}; \mathcal{K}) = \mathcal{H}_+^2(\mathbb{R}; \mathcal{K}) \oplus \mathcal{H}_-^2(\mathbb{R}; \mathcal{K})$. We denote the orthogonal projections in $L^2(\mathbb{R}; \mathcal{K})$ on $\mathcal{H}_+^2(\mathbb{R}; \mathcal{K})$ and $\mathcal{H}_-^2(\mathbb{R}; \mathcal{K})$, respectively, by P_+ and P_- . Let $L^2(\mathbb{R}^\pm; \mathcal{K})$ be the subspaces of $L^2(\mathbb{R}; \mathcal{K})$ consisting of square-integrable functions supported on $\mathbb{R}^\pm$. There exists another orthogonal decomposition $L^2(\mathbb{R}; \mathcal{K}) = L^2(\mathbb{R}^-; \mathcal{K}) \oplus L^2(\mathbb{R}^+; \mathcal{K})$ with projections on $P_{\mathbb{R}^+}$ and $P_{\mathbb{R}^-}$. The following theorem is proved in [7]:

Theorem II.2 *Let $M : L^2(\mathbb{R}^+; \mathcal{K}) \mapsto L^2(\mathbb{R}^+; \mathcal{K})$ be the operator defined by*

$$M := (P_{\mathbb{R}^+} P_+ P_{\mathbb{R}^+})|_{L^2(\mathbb{R}^+; \mathcal{K})}. \quad (3)$$

Then M is a positive, contractive and injective operator on $L^2(\mathbb{R}^+; \mathcal{K})$, such that $\text{Ran } M$ is dense in $L^2(\mathbb{R}^+; \mathcal{K})$ and M is a forward Lyapunov operator for the evolution on $L^2(\mathbb{R}^+; \mathcal{K})$ defined in Eq. (2), i.e., for every $\psi \in L^2(\mathbb{R}^+; \mathcal{K})$ we have

$$(\psi(t_2), M\psi(t_2)) \leq (\psi(t_1), M\psi(t_1)), \quad t_2 \geq t_1 \geq 0, \quad \psi(t) = U(t)\psi, \quad (4)$$

and moreover $\lim_{t \rightarrow \infty} (\psi(t), M\psi(t)) = 0$. □

The following corollary to Theorem II.2 casts Eq. (3) in a more useful form:

Corollary II.3 *Let M be the Lyapunov operator given in Eq. (3). Then, for any function $f \in L^2(\mathbb{R}^+; \mathcal{K})$ we have*

$$(Mf)(E) = -\frac{1}{2\pi i} \int_0^\infty dE' \frac{1}{E - E' + i0^+} f(E'), \quad E \in \mathbb{R}^+, \quad f \in L^2(\mathbb{R}^+; \mathcal{K}). \quad (5)$$

□

Proof: For every function $g \in \mathcal{H}^2(\mathbb{C}^\pm; \mathcal{K})$ one has [8]

$$\mp \frac{1}{2\pi i} \int_{-\infty}^\infty dt \frac{1}{z - t} g_\pm(t) = \begin{cases} g(z), & \pm \text{Im } z > 0 \\ 0, & \pm \text{Im } z < 0 \end{cases}, \quad (6)$$

where $g_\pm \in \mathcal{H}_\pm^2(\mathbb{R}; \mathcal{K})$ is the boundary value function of g on $\mathbb{R}$. Hence, the orthogonal projection P_+ of $L^2(\mathbb{R}^+; \mathcal{K})$ on $\mathcal{H}_+^2(\mathbb{R}; \mathcal{K})$ is explicitly given by

$$(P_+g)(\sigma) = -\frac{1}{2\pi i} \int_{-\infty}^\infty d\sigma' \frac{1}{\sigma - \sigma' + i0^+} g(\sigma'), \quad g \in L^2(\mathbb{R}; \mathcal{K}), \quad (7)$$

where Eq. (7) hold a.e. for $\sigma \in \mathbb{R}$. From Eqns. (3), (7) we then immediately obtain Eq. (5). ■

Theorem II.4 *The spectrum of M satisfies $\sigma(M) = \sigma_{ac}(M) = [0, 1]$. In $L^2(\mathbb{R}^+; \mathcal{K})$ choose an orthogonal basis $\{\mathbf{e}_\lambda(E)\}_{E \in \mathbb{R}^+; \lambda \in \Lambda}$ with Λ an appropriate index set such that for every $E \in \mathbb{R}^+$ the set $\{\mathbf{e}_\lambda(E)\}_{\lambda \in \Lambda}$ is a basis of $\mathcal{K}$ and we have $(\mathbf{e}_\lambda(E), \mathbf{e}_{\lambda'}(E'))_{L^2(\mathbb{R}^+; \mathcal{K})} = \delta(E - E')\delta_{\lambda\lambda'}$, where $\delta_{\lambda\lambda'}$ stands for the Kronecker delta for discrete indices and the Dirac delta for continuous indices. Then for $m \in (0, 1)$ the function defined by*

$$\mathbf{g}_{m,\lambda} = \int_0^\infty dE \frac{E^{-\frac{i}{2\pi} \ln(\frac{1-m}{m}) - \frac{1}{2}}}{2\pi\sqrt{m(1-m)}} \mathbf{e}_\lambda(E) \quad (8)$$

is a generalized eigenfunction of M satisfying $M \mathbf{g}_{m,\lambda} = m \mathbf{g}_{m,\lambda}$. These eigenfunctions are normalized in such a way that $(\mathbf{g}_{m,\lambda}, \mathbf{g}_{m',\lambda'})_{L^2(\mathbb{R}^+; \mathcal{K})} = \delta(m - m')\delta_{\lambda\lambda'}$, and we have the eigenfunction expansion

$$M = \sum_{\lambda \in \Lambda} \int_0^1 dm \, m \, \mathbf{g}_{m,\lambda} \mathbf{g}_{m,\lambda}^*. \quad (9)$$

□

Proof: Consider the eigenvalue equation $M \mathbf{g}_{m,\lambda} = m \mathbf{g}_{m,\lambda}$. In the basis $\{\mathbf{e}_\lambda(E)\}_{E \in \mathbb{R}^+; \lambda \in \Lambda}$ the kernel of M is given by $(\mathbf{e}_\lambda(E), M \mathbf{e}_{\lambda'}(E')) = -(2\pi i)^{-1}(E - E' + i0^+)^{-1} \delta_{\lambda\lambda'}$. Hence, the eigenvalue equation for M assumes the form

$$-\frac{1}{2\pi i} \int_0^\infty dE' \frac{1}{E - E' + i0^+} g_{m,\lambda}(E') = m g_{m,\lambda}(E), \quad E \in \mathbb{R}^+. \quad (10)$$

where $g_{m,\lambda}(E) = (\mathbf{e}_\lambda(E), \mathbf{g}_{m,\lambda})$. Assume that $m \in (0, 1)$. To solve Eq. (10) we assume that the function $g_{m,\lambda}(E)$ is the boundary value on $\mathbb{R}^+$ from above of an analytic function $\tilde{g}_{m,\lambda}(z)$ defined in $\mathbb{C} \setminus \mathbb{R}^+$. So that $g_{m,\lambda}(E) = \tilde{g}_{m,\lambda}(E + i0^+)$. This allows us to analytically continue eq. (10) into the cut complex plane

$$-\frac{1}{2\pi i} \int_0^\infty dE' \frac{1}{z - E'} g_{m,\lambda}(E') = m \tilde{g}_{m,\lambda}(z), \quad \text{Im} z \neq 0. \quad (11)$$

Taking in Eq. (11) the difference between the limits from above and below $\mathbb{R}^+$ we get

$$g_{m,\lambda}(E) = \tilde{g}_{m,\lambda}(E + i0^+) = m (\tilde{g}_{m,\lambda}(E + i0^+) - \tilde{g}_{m,\lambda}(E - i0^+)). \quad (12)$$

The function $\tilde{g}_{m,\lambda}(z)$ can now be continued to a second Riemann sheet by making use of the branch cut along $[0, \infty)$ in eq. (11). Denoting this two sheeted function again by $\tilde{g}_{m,\lambda}(z)$, Eq. (11), reduces to

$$\tilde{g}_{m,\lambda}(e^{2\pi i} z) = -\frac{1-m}{m} \tilde{g}_{m,\lambda}(z). \quad (13)$$

Eq. (13) admits solutions of the form $\tilde{g}_{m,\lambda}(z) = N_m z^\beta$ with $\beta = k + \frac{1}{2} - \frac{i}{2\pi} \ln\left(\frac{1-m}{m}\right)$, $k \in \mathbb{Z}$, and N_m a normalization factor dependent on m . Setting $k = -1$ and $N_m = (4\pi^2 m (1-m))^{-1/2}$ the solutions of the eigenvalue equation, Eq. (10), satisfy $\int_0^\infty dE \overline{g_{m',\lambda'}(E)} g_{m,\lambda}(E) = \delta(m - m') \delta_{\lambda\lambda'}$. With this choice of k and N_m the generalized eigenfunctions of M are given by Eq. (8).

We proceed to show by construction that the set of generalized eigenfunctions, Eq. (8), forms a complete set for the operator M . Consider the following integral

$$\sum_\lambda \int_0^1 dm m \mathbf{g}_{m,\lambda} \mathbf{g}_{m,\lambda}^* = \sum_\lambda \int_0^\infty dE \int_0^\infty dE' \frac{\mathbf{e}_\lambda(E) \mathbf{e}_\lambda^*(E')}{4\pi^2 \sqrt{E E'}} \int_0^1 \frac{dm}{1-m} \left(\frac{m}{1-m} \right)^{\frac{i}{2\pi} \ln(E/E')}. \quad (14)$$

where dm is a Lebesgue measure on $[0, 1]$. The right hand side of eq. (14) is readily obtained from that of eq. (8). Recall the definition of the Euler beta function $B(x, y) = \int_0^1 dm m^{x-1} (1-m)^{y-1} = \Gamma(x) \Gamma(y) \Gamma^{-1}(x+y)$. $B(x, y)$ is well defined for $\text{Re } x > 0$ and $\text{Re } y > 0$, and can be analytically continued to other parts of the complex x and y planes. However, it is not well defined for $x = 0$ and $y = 0$, i.e. whenever $E = E'$ in Eq. (14). To avoid this problem we shift E (or E') away from the real axis. Hence, in the integration over m on the right hand side of Eq. (14) we take the limit

$$\begin{aligned} & \frac{1}{4\pi^2} (E E')^{-1/2} \lim_{\theta \rightarrow 0^+} \int_0^1 dm (1-m)^{-\frac{i}{2\pi} \ln(e^{i\theta} E/E') - 1} m^{\frac{i}{2\pi} \ln(e^{i\theta} E/E')} \\ &= \frac{1}{4\pi^2} (E E')^{-1/2} \lim_{\theta \rightarrow 0^+} B\left(1 - \frac{\theta}{2\pi} + \frac{i}{2\pi} \ln\left(\frac{E}{E'}\right), \frac{\theta}{2\pi} - \frac{i}{2\pi} \ln\left(\frac{E}{E'}\right)\right) \\ &= \frac{1}{4\pi^2} (E E')^{-1/2} \lim_{\theta \rightarrow 0^+} \frac{\pi}{\sin\left(-\frac{i}{2\pi} \ln\left(\frac{e^{i\theta} E}{E'}\right)\right)} = -\frac{1}{2\pi i} \frac{1}{E - E' + i0^+}. \end{aligned} \quad (15)$$

where in the last line in Eq. (15) we used the identity $\Gamma(z) \Gamma(1-z) = \pi \sin^{-1}(\pi z)$. Thus, the intgral on the spectrum of M on the left hand side of Eq. (15) reconstructs the kernel of M in Eq. (5), i.e., we have reconstructed the operator M using the set of generalized eigenfunctions $\{\mathbf{g}_{m,\lambda}\}_{m \in (0,1), \lambda \in \Lambda}$. Taken together with Eq. (14), Eq. (15) also shows that M has no point spectrum at $m = 0$ and $m = 1$ and that $\sigma(M) = \sigma_{ac}(M) = [0, 1]$. ■

Consider a quantum mechanical problem for which the Hamiltonian H , defined on an appropriate separable Hilbert space $\mathcal{H}$, satisfies the conditions: (i) Its ac spectrum is $\sigma_{ac}(H) = \mathbb{R}^+$. (ii) The multiplicity of $\sigma_{ac}(H)$ is uniform on $\mathbb{R}^+$. Let $\mathcal{H}_{ac}$ be the subspace of $\mathcal{H}$ corresponding to the ac spectrum of H . Then $\mathcal{H}_{ac}$ has a representation in terms of a function space $L^2(\mathbb{R}^+; \mathcal{K})$, where $\mathcal{K}$ is a Hilbert space whose dimension correpsonds to the multiplicity of $\sigma_{ac}(H)$ and the evolution generated by H is represented by $\{U(t)\}_{t \in \mathbb{R}}$ defined in Eq. (2). For such models we can apply the results discussed above. Using the standard Dirac notation, Eq. (2) for M can be rewritten as




Figure 1: Monotonic decrease of the expectation value of M for a free Gaussian wave-packet with $p_0 = 0.64\eta$ and $\xi_0 = 0.3\eta$.

$$M = -\frac{1}{2\pi i} \sum_{\lambda} \int_0^{\infty} dE \int_0^{\infty} dE' |E, \lambda\rangle \frac{1}{E - E' + i0^+} \langle E', \lambda|, \quad (16)$$

where the summation over λ stands for summation over discrete degeneracy indices and appropriate integration over continuous degeneracy indices.

As an example we take a one-dimensional Gaussian wave-packet representing the propagation free particle of mass η along $\mathbb{R}^+$

$$\psi(x, t) = \left(\frac{\eta^2 \xi_0^2}{\pi (\eta + i\xi_0^2 t)^2} \right)^{1/4} \exp \left(-\frac{\eta \xi_0^2 x^2 + ip_0 (p_0 t - 2\eta x)}{2(\eta + i\xi_0^2 t)} \right). \quad (17)$$

p_0 and ξ_0 are the location and width of the wave-packet in momentum space at $t = 0$. After obtaining the generalized eigenfunctions of the free Hamiltonian, constructing the Lyapunov operator M according to Eqs. (16) and (14), and applying it to the wave-packet in Eq. (17) we obtain Fig. 1, showing the time evolution of the expectation value of M , and Fig. 2, showing the time evolution of the associated probability density function in the position basis and that of the eigenstates of M . We see that if the sequence of time frames is shown in reverse order one is unable to tell whether time is running backwards or whether one is observing a Gaussian wave-packet propagating along the $\mathbb{R}^-$ (with time running forward). However, if to all frames we attach the expectation value of M , then it possible to distinguish between these two scenarios. The example plainly illustrates the time-ordering of states introduced into the Hilbert space by the existence of a Lyapunov operator.




Figure 2: Time frames of $|\psi(x, t)|^2$ and $|\langle m | \psi(t) \rangle|^2$ for the Gaussian wave-packet of Fig. 1.

In a forthcoming publication it will be shown that the existence of the operator M leads to an irreversible representation of the quantum dynamics, and to a novel representation of the scattering process, in which the contribution of the resonance is singled out.

Addendum – This paper presents a mathematically rigorous reworking of some of the results appearing in [13], in particular, the derivation of the operator's spectrum and generalized eigenfunctions. The operator has since been generalized to a larger class of self-adjoint Lyapunov operators [14].

Acknowledgements

Y.S. acknowledges the support of the Israeli Science Foundation Grant no. 1169/06, and the Center for Advanced Studies in Mathematics at Ben-Gurion University, where part of this research was conducted. J.S. and S.M. acknowledge the support of the Israeli Science Foundation Grant no. 784/06.

-
- [1] See, for example, G.W. Mackey, *The Theory of Unitary Group Representations*, University of Chicago Press, 1976.
 - [2] W.E. Pauli, in *Handbuch der Physik*, Springer-Verlag, 1926.
 - [3] E.A. Galapon, Pauli's theorem and quantum canonical pairs: the consistency of a bounded, self adjoint time operator canonically conjugate to a Hamiltonian with nonempty point spectrum, *Proc. R. Soc. Lond. A* (2002) 451-472.
 - [4] W.G. Unruh, R.M. Wald, Time and the interpretation of canonical quantum gravity, *Phys. Rev. D* 40 (1989) 2598-2614.
 - [5] B. Misra, I. Prigogine, M. Courbage, Lyapounov variable: Entropy and measurement in quantum mechanics, *Proc. Natl. Acad. Sci. USA* 76 (1979) 4768-4772.
 - [6] A.S. Holevo, *Probabilistic and Statistical Aspects of Quantum Theory*, North-Holland, 1982.
 - [7] Y. Strauss, Self-adjoint Lyapunov variables, temporal ordering and irreversible representations of Schrödinger evolution, *J. Math. Phys.* 51 (2010) 022104. See also [13].
 - [8] R.E.A.C. Paley, N. Wiener, in: *Fourier Transforms in the Complex Domain*, American Mathematical Society, 1934.
 - [9] See, for example, H.M. Evans, *Perturbation Theory and Atomic Resonances Since Schrödinger's Time*, in: F. Gesztes, P. Deift, C. Galvez, P. Perry, W. Schlag (Eds.), *Spectral Theory and Mathematical Physics: A Festschrift in Honor of Barry Simon's 60th Birthday*, American Mathematical Society, 2007.
 - [10] L.P. Horwitz, J.-P. Marchand, The decay-scattering system, *Rocky Mt. J. Math.* 1 (1971) 225-254.
 - [11] Y. Strauss, On the semigroup decomposition of the time evolution of quantum mechanical resonances, *J. Math. Phys.* 46 (2005) 102109.
 - [12] Y. Strauss, L.P. Horwitz, A. Volovick, Approximate resonance states in the semigroup decomposition of resonance evolution, *J. Math. Phys.* 47 (2006) 123505.
 - [13] Y. Strauss, J. Silman, S. Machnes, L.P. Horwitz, arXiv:0802.2448 [quant-ph].
 - [14] G.C. Hegerfeldt, J.G. Muga, J. Muñoz, *Phys. Rev. A* 82 (2010) 012113.